

The geometry of dissipation

Asier López Gordón

Instituto de Ciencias Matemáticas (ICMAT)

Supervisor: Manuel de León Rodríguez

Tutor: Rafael Orive Illera

Programa de Doctorado en Matemáticas

Universidad Autónoma de Madrid



Geometric frameworks for dynamics

- By the Equivalence Principle, the laws of physics are the same for all observers, i.e., for all systems of coordinates.
- Hence, we should formulate the laws of physics in a coordinate-independent language, namely, the language of differential geometry.

Geometric frameworks for Hamiltonian dynamics

- As it is well-known, symplectic manifolds are the natural framework for Hamiltonian mechanics.
- Hamiltonian dynamics are conservative: the Hamiltonian flow preserves the symplectic form and the Hamiltonian function.

Geometric frameworks for dissipative dynamics

- The ubiquity of physical phenomena where the energy or the volume of the phase space are not preserved leads to the necessity of developing frameworks for non-conservative dynamics.
- In this dissertation, we consider three geometric frameworks for dissipative dynamics:
 - contact Hamiltonian (and Lagrangian) systems,
 - Hamiltonian (and Lagrangian) systems with external forces,
 - mechanical systems with impacts.
- We have generalized several results from conservative systems to these frameworks. In particular, those results concerning symmetries, reduction and integrability.

Contact geometry

Definition

A (co-oriented) **contact manifold** is a pair (M, η) , where M is an $(2n + 1)$ -dimensional manifold and η is a 1-form on M such that the map

$$\begin{aligned} b_\eta: \mathfrak{X}(M) &\rightarrow \Omega^1(M) \\ X &\mapsto \iota_X d\eta + \eta(X)\eta, \end{aligned}$$

is an isomorphism of $\mathcal{C}^\infty(M)$ -modules.

- There exists a unique vector field R on (M, η) , called the **Reeb vector field**, given by $R = b_\eta^{-1}(\eta)$, or, equivalently,

$$\iota_R d\eta = 0, \quad \iota_R \eta = 1.$$

Contact geometry

- The **Hamiltonian vector field** of $f \in \mathcal{C}^\infty(M)$ is given by

$$X_f = b_\eta^{-1}(df) - (R(f) + f)R,$$

- Around each point on M there exist **Darboux coordinates** (q^i, p_i, z) such that

$$\eta = dz - p_i dq^i,$$

$$R = \frac{\partial}{\partial z},$$

$$X_f = \frac{\partial f}{\partial p_i} \frac{\partial}{\partial q^i} - \left(\frac{\partial f}{\partial q^i} + p_i \frac{\partial f}{\partial z} \right) \frac{\partial}{\partial p_i} + \left(p_i \frac{\partial f}{\partial p_i} - f \right) \frac{\partial}{\partial z}.$$

Contact Hamiltonian systems

Definition

A **contact Hamiltonian system** is a triple (M, η, h) formed by a contact manifold (M, η) and a **Hamiltonian function** $h \in \mathcal{C}^\infty(M)$.

- The dynamics of (M, η, h) is determined by the integral curves of the Hamiltonian vector field X_h of h w.r.t. η .

Contact Hamiltonian systems

- In Darboux coordinates, these curves $c(t) = (q^i(t), p_i(t), z(t))$ are determined by the **contact Hamilton equations**:

$$\frac{dq^i(t)}{dt} = \frac{\partial h}{\partial p_i} \circ c(t),$$

$$\frac{dp_i(t)}{dt} = -\frac{\partial h}{\partial q^i} \circ c(t) + p_i(t) \frac{\partial h}{\partial z} \circ c(t),$$

$$\frac{dz(t)}{dt} = p_i(t) \frac{\partial h}{\partial p_i} \circ c(t) - h \circ c(t).$$

Contact Lagrangian systems

- The Darboux coordinate z can be regarded, from a variational point of view, as the action functional.
- Let L be an **action-dependent** Lagrangian function.
- If L is regular, the Legendre transformation leads to a contact Hamiltonian system.
- Very loosely, the Herglotz functional \mathcal{A} is like the usual action functional, but instead of being given by an integral is given by the ODE

$$\frac{d}{dt}\mathcal{A}[q(t)] = L\left(q(t), \dot{q}(t), \mathcal{A}[q(t)]\right).$$

- One seeks for curves $q: I \subseteq \mathbb{R} \rightarrow Q$ that are extremals of \mathcal{A} .

Contact Lagrangian systems

- Given two fixed points $q_1, q_2 \in Q$ and an interval $[a, b]$, let

$$\Omega(q_1, q_2, [a, b]) = \left\{ c \in \mathcal{C}^2([a, b] \rightarrow Q) \mid c(a) = q_1, c(b) = q_2 \right\}.$$

- Consider the operator

$$\mathcal{Z}: \Omega(q_1, q_2, [a, b]) \rightarrow \mathcal{C}^2([a, b] \rightarrow \mathbb{R})$$

that assigns to each curve c the solution $\mathcal{Z}(c)$ of the following Cauchy problem:

$$\begin{aligned} \frac{d\mathcal{Z}(c)(t)}{dt} &= L(c(t), \dot{c}(t), \mathcal{Z}(c)(t)), \\ \mathcal{Z}(c)(a) &= z_a. \end{aligned}$$

Contact Lagrangian systems

- The **Herglotz action functional** is the map

$$\begin{aligned} \mathcal{A}: \Omega(q_1, q_2, [a, b]) &\rightarrow \mathbb{R} \\ c &\mapsto \mathcal{Z}(c)(b). \end{aligned}$$

- A curve $c \in \Omega(q_1, q_2, [a, b])$ is a critical point of \mathcal{A} (i.e., $d\mathcal{A}(c) = 0$) if and only if it satisfies the **Herglotz–Euler–Lagrange equations**:

$$\begin{aligned} \frac{d}{dt} \frac{\partial L}{\partial v^i} \circ \chi(t) - \frac{\partial L}{\partial q^i} \circ \chi(t) - \frac{\partial L}{\partial v^i} \circ \chi(t) \frac{\partial L}{\partial z} \circ \chi(t) &= 0, \\ \frac{dz}{dt} &= L \circ \chi(t), \end{aligned}$$

where $\chi(t) = (c(t), \dot{c}(t), z(t))$.

Dissipated quantities

- In contact Hamiltonian dynamics dissipated quantities are akin to conserved quantities in symplectic dynamics.
- Energy (Hamiltonian function) is no longer conserved, but dissipated in a certain manner:

$$X_h(h) = -R(h)h.$$

Dissipated quantities

Example (linear dissipation)

Let $M = \mathbb{R}^3$ with canonical coordinates (q, p, z) ,

$$\eta = dz - pdq, \quad h = \frac{p^2}{2} + V(q) + \kappa z, \quad \kappa \in \mathbb{R}.$$

Then $X_h(h) = -\kappa h$, so

$$h \circ c(t) = e^{-\kappa t} h \circ c(0),$$

along an integral curve c of X_h .

Dissipated quantities

Definition

Let (M, η, h) be a contact Hamiltonian system. A **dissipated quantity** is a solution $f \in \mathcal{C}^\infty(M)$ to the PDE

$$X_h(f) = -R(h)f.$$

Jacobi structure of a contact manifold

- The **Jacobi bracket** $\{\cdot, \cdot\}: \mathcal{C}^\infty(M) \times \mathcal{C}^\infty(M) \rightarrow \mathcal{C}^\infty(M)$ is given by

$$\{f, g\} = -d\eta(b_\eta^{-1}df, b_\eta^{-1}dg) - fR(g) + gR(f).$$

- It is a Lie bracket, namely, it is bilinear, skew-symmetric and satisfies the Jacobi identity.
- It satisfies the weak Leibniz identity:

$$\{f, gh\} = \{f, g\}h + \{f, h\}g - ghR(f).$$

Jacobi brackets and dissipated quantities

- The Jacobi bracket can also be expressed as follows:

$$\{f, g\} = X_f(g) + gR(f).$$

Proposition

Let (M, η, h) be a contact Hamiltonian system and let $f \in \mathcal{C}^\infty(M)$. Then, f is a dissipated quantity (i.e., $X_h(f) = -R(h)f$) iff

$$\{f, h\} = 0.$$

Functions in involution

Definition

Let $\{\cdot, \cdot\}$ be a Jacobi bracket on M . A collection of functions $f_1, \dots, f_k \in \mathcal{C}^\infty(M)$ are said to be **in involution** if

$$\{f_i, f_j\} = 0, \quad \forall i, j \in \{1, \dots, k\}.$$

Remark

Unlike in the case of Poisson brackets, f_i and f_j being in involution does not imply that X_{f_i} is tangent to the level sets of f_j . Consequently, the submanifolds

$$M_\Lambda = \bigcap_{i=1}^k f_i^{-1}(\Lambda_i), \quad \Lambda_i \in \mathbb{R}$$

are no longer invariant under the flows of X_{f_1}, \dots, X_{f_k} .

Liouville–Arnol'd theorem for contact Hamiltonian systems

- **Crucial idea:** replace the level sets (i.e., preimages of points) M_Λ by **preimages of rays**

$$M_{\langle \Lambda \rangle_+} = \{x \in M \mid \exists r \in \mathbb{R}^+ : f_\alpha(x) = r\Lambda_\alpha \forall \alpha\},$$

with $\alpha \in \{0, 1, \dots, n\}$ and $\Lambda = (\Lambda_0, \Lambda_1, \dots, \Lambda_n) \in \mathbb{R}^{n+1} \setminus \{0\}$.

Liouville–Arnol'd theorem for contact Hamiltonian systems

Theorem (Colombo, de León, Lainz, L. G., 2023)

Let (M, η) be a $(2n + 1)$ -dimensional contact manifold. Suppose that f_0, f_1, \dots, f_n are functions in involution such that $\text{rank}\{df_\alpha\}_\alpha \geq n$. Then, $M_{\langle \Lambda \rangle_+}$ is invariant by the Hamiltonian flow of f_α and diffeomorphic to $\mathbb{T}^k \times \mathbb{R}^{n+1-k}$.

Moreover, there is a neighborhood U of $M_{\langle \Lambda \rangle_+}$ such that

- 1 There exists coordinates $(y^0, \dots, y^n, \tilde{A}_1, \dots, \tilde{A}_n)$ on U in which the equations of motion read

$$\dot{y}^\alpha = \Omega^\alpha(\tilde{A}_i), \quad \dot{\tilde{A}}_i = 0, \quad \alpha \in \{0, \dots, n\}, \quad i \in \{1, \dots, n\}.$$

- 2 There exists a conformal change $\tilde{\eta} = \eta/A_0$ such that (y^i, \tilde{A}_i, y^0) are Darboux coordinates for $(M, \tilde{\eta})$, i.e. $\tilde{\eta} = dy^0 - \tilde{A}_i dy^i$.

Steps of the proof

- 1 Symplectize (M, η) and f_α , obtaining an exact symplectic manifold (M^Σ, θ) and homogeneous functions in involution f_α^Σ .
- 2 Prove a Liouville–Arnold theorem for exact symplectic manifolds with homogeneous functions in involution.
- 3 “Un-symplectize” the action-angle coordinates $(y_\Sigma^\alpha, A_\Sigma^\alpha)$ on M^Σ , yielding functions (y^α, A_Σ) on M .
- 4 Introduce action-angle coordinates (y^α, \tilde{A}_i) on M , where $\tilde{A}_i = -\frac{A_i}{A_0}$.

Example

- Let $M = \mathbb{R}^3 \setminus \{0\}$ with canonical coordinates (q, p, z) , and $\eta = dz - pdq$.
- The functions $h = p$ and $f = z$ are in involution.
- We have a chart $(M \setminus \{z = 0\}; y^0, y^1, \tilde{A})$, where

$$y^0 = q, \quad y^1 = -\log z \quad \tilde{A} = -\frac{p}{z}.$$

- In this chart,

$$X_h = \frac{\partial}{\partial y^0}, \quad X_f = \frac{\partial}{\partial y^1}$$

- It is a Darboux chart for the contact form

$$\tilde{\eta} = \frac{1}{A_0} \eta = dy^0 - \tilde{A} dy^1.$$

Dissipated quantities and stability

In a work in progress, we employ dissipated quantities to study the stability of contact Hamiltonian systems.

Proposition (de Lucas, L. G., Zawora)

Let (M, η, h) be a contact Hamiltonian system such that $X_h(x_0) = 0$. Suppose that f_1, \dots, f_k are dissipated quantities. If $(Rh)(x_0) > 0$ at an isolated point $x_0 \in \bigcap_{i=1}^k f_i^{-1}(0)$, then x_0 is an asymptotically stable equilibrium point.

Cocontact structures

- We would like to model dynamical systems which are dissipative and have an explicit time dependence.
- For instance, consider a friction force that changes with time.
- Idea: adding explicit time dependence to contact dynamics.

Cocontact structures

Definition (de León, Gaset, Gràcia, Muñoz-Lecanda, and Rivas, 2022)

A **cocontact manifold** is a triple (M, τ, η) such that:

- 1 M is a $(2n + 2)$ -dimensional manifold,
- 2 τ and η are 1-forms,
- 3 $d\tau = 0$,
- 4 The map

$$b_{(\tau, \eta)}: \mathfrak{X}(M) \rightarrow \Omega^1(M)$$
$$X \mapsto (\iota_X \tau)\tau + \iota_X d\eta + (\iota_X \eta)\eta$$

is an isomorphism of $\mathcal{C}^\infty(M)$ -modules.

Reeb and Hamiltonian vector fields

- **Reeb vector fields:** $R_t = b_{(\tau, \eta)}^{-1}(\tau)$, $R_z = b_{(\tau, \eta)}^{-1}(\eta)$.
- **Hamiltonian vector field:**

$$X_f = b_{(\tau, \eta)}^{-1}(df) - (R_z(f) + f) R_z + (1 - R_t(f)) R_t.$$

- **Darboux coordinates** (t, q^i, p_i, z) :

$$\tau = dt, \quad \eta = dz - p_i dq^i, \quad R_t = \frac{\partial}{\partial t}, \quad R_z = \frac{\partial}{\partial z},$$

$$X_f = \frac{\partial}{\partial t} + \frac{\partial f}{\partial p_i} \frac{\partial}{\partial q^i} - \left(\frac{\partial f}{\partial q^i} + p_i \frac{\partial f}{\partial z} \right) \frac{\partial}{\partial p_i} + \left(p_i \frac{\partial f}{\partial p_i} - f \right) \frac{\partial}{\partial z}.$$

Dissipation of energy

Definition

A **cocontact Hamiltonian system** is a tuple (M, τ, η, h) formed by a cocontact manifold (M, τ, η) and a **Hamiltonian function** $h \in \mathcal{C}^\infty(M)$.

The energy of a cocontact Hamiltonian system is not preserved due to both the “contact variable” and the time dependence. Indeed,

$$X_h(h) = -R_z(h)h + R_t(h).$$

Dissipated quantities

Definition

Let (M, τ, η, h) be a cocontact Hamiltonian system. A **dissipated quantity** is a function $f: M \rightarrow \mathbb{R}$ such that

$$X_h(f) = -R_z(h)f.$$

Noether's theorem for cocontact Hamiltonian systems

Theorem (Gaset, L. G., Rivas, 2023)


Consider the cocontact Hamiltonian system (M, τ, η, h) . Let $Y \in \mathfrak{X}(M)$.

- 1 If $\eta([Y, X_h]) = 0$ and $\tau(Y) = 0$, then $f = -\eta(Y)$ is a dissipated quantity.
- 2 Conversely, given a dissipated quantity f , the vector field $Y = X_f - R_t$ verifies $\eta([Y, X_h]) = 0$, $\tau(Y) = 0$ and $f = -\eta(Y)$.

Definition

A **generalized infinitesimal dynamical symmetry** is a vector field $Y \in \mathfrak{X}(M)$ such that $\eta([Y, X_h]) = 0$ and $\tau(Y) = 0$.

Classification of infinitesimal symmetries



Example (The two-body problem with time-dependent friction)

- The phase space is $\mathbb{R} \times T^*\mathbb{R}^6 \times \mathbb{R}$, with coords. $(t, \mathbf{q}^1, \mathbf{q}^2, \mathbf{p}_1, \mathbf{p}_2, z)$, where $\mathbf{q}^a \in \mathbb{R}^3$ is the position of the body $a \in \{1, 2\}$ and $\mathbf{p}_a \in \mathbb{R}^3$ is its momentum.
- The Hamiltonian function is

$$H = \frac{\|\mathbf{p}_1\|^2}{2m_1} + \frac{\|\mathbf{p}_2\|^2}{2m_2} + U(\|\mathbf{q}^2 - \mathbf{q}^1\|) + \gamma(t)z,$$

and the cocontact structure is given by the one-forms

$$\eta = dz - \mathbf{p}_1 \cdot d\mathbf{q}^1 - \mathbf{p}_2 \cdot d\mathbf{q}^2, \quad \tau = dt.$$

- The vector fields $Y_i = \frac{\partial}{\partial q_i^1} + \frac{\partial}{\partial q_i^2}$, $i \in \{1, 2, 3\}$ are infinitesimal strict Hamiltonian symmetries and the associated dissipated quantities are the components of $\mathbf{p}_1 + \mathbf{p}_2$.

Section 3

Systems with external forces

Forced Hamiltonian systems

Given a manifold Q , let T^*Q be its cotangent bundle with canonical one-form θ_Q and canonical symplectic form $\omega_Q = -d\theta_Q$.

Definition

A **forced Hamiltonian system** is a triple (Q, h, α) where Q is a manifold, $h \in \mathcal{C}^\infty(T^*Q)$ is a function and $\alpha \in \Omega^1(T^*Q)$ is a semibasic one-form (i.e., $\alpha(X) = 0$ for any vertical vector field X). The **forced Hamiltonian vector field** $X_{h,\alpha} \in \mathfrak{X}(T^*Q)$ is given by

$$\iota_{X_{h,\alpha}} \omega_Q = dh + \alpha.$$

Local expressions

Given local coordinates (q^i) on Q and the induced bundle coordinates (q^i, p_i) on T^*Q , we have that

$$\theta_Q = p_i dq^i,$$

$$\omega_Q = dq^i \wedge dp_i,$$

$$\alpha = \alpha_i(q, p) dq^i,$$

$$X_{h, \alpha} = \frac{\partial h}{\partial p_i} \frac{\partial}{\partial q^i} - \left(\frac{\partial h}{\partial q^i} + \alpha_i \right) \frac{\partial}{\partial p_i}.$$

Reduction *à la* Marsden–Weinstein

- Let G be a Lie group with Lie algebra \mathfrak{g} , and dual \mathfrak{g}^* .
- Consider a Lie group action $\Phi: G \times Q \rightarrow Q$ of G on a manifold Q and its cotangent lift $\Phi^{T^*}: G \times T^*Q \rightarrow T^*Q$.
- Henceforth, assume that both of these actions are free and proper.

Reduction *à la* Marsden–Weinstein

- Let $\xi_{T^*Q} \in \mathfrak{X}(T^*Q)$ denote the infinitesimal generator of the action defined by $\xi \in \mathfrak{g}$, i.e.,

$$\xi_{T^*Q}(x) = \left. \frac{d}{dt} \right|_{t=0} \Phi_{\exp t\xi}^{T^*}(x),$$

where $\exp: \mathfrak{g} \rightarrow G$ is the exponential map.

- Let $\mathbf{J}: T^*Q \rightarrow \mathfrak{g}^*$ denote the natural momentum map, namely,

$$\mathbf{J}^\xi(x) := \langle \mathbf{J}(x), \xi \rangle = \iota_{\xi_{T^*Q}} \theta_Q(x),$$

Reduction à la Marsden–Weinstein

Proposition

Let (Q, h, α) be a forced Hamiltonian system such that h is Φ^{T^*} -invariant.

- ① For each $\xi \in \mathfrak{g}$, the function J^ξ is a conserved quantity iff

$$\alpha(\xi_{T^*Q}) = 0. \quad (1)$$

- ② If Eq. (1) holds, then α is ξ -invariant iff

$$\iota_{\xi_{T^*Q}} d\alpha = 0.$$

- ③ The subset

$$\mathfrak{g}_\alpha = \{ \xi \in \mathfrak{g} \mid \alpha(\xi_{T^*Q}) = 0, \iota_{\xi_{T^*Q}} d\alpha = 0 \}$$

is a Lie subalgebra of \mathfrak{g} .

Reduction *à la* Marsden–Weinstein

- Let G_α be the unique connected Lie subgroup of G whose Lie algebra is \mathfrak{g}_α .
- Assume that G_α is a closed Lie subgroup of G .
- Let μ be a regular value of the natural momentum map $\mathbf{J}_\alpha: T^*Q \rightarrow \mathfrak{g}_\alpha^*$.
- Denote by $G_{\alpha,\mu} \subseteq G_\alpha$ the isotropy subgroup of μ w.r.t. the coadjoint action.

Reduction à la Marsden–Weinstein

Theorem (de León, Lainz, L. G., 2021)

- 1 $X_{h,\alpha}$ is tangent to $\mathbf{J}_\alpha^{-1}(\mu)$.
- 2 $M_\mu := \mathbf{J}_\alpha^{-1}(\mu)/G_{\alpha,\mu}$ has a symplectic form ω_μ uniquely determined by

$$\pi_\mu^* \omega_\mu = i_\mu^* \omega_Q,$$

where the maps $i_\mu: \mathbf{J}_\alpha^{-1}(\mu) \hookrightarrow \mathbf{T}^*Q$ and $\pi_\mu: \mathbf{J}_\alpha^{-1}(\mu) \rightarrow \mathbf{J}_\alpha^{-1}(\mu)/G_{\alpha,\mu}$ denote the inclusion and the projection, respectively.

- 3 We have a reduced Hamiltonian function h_μ and force α_μ on M_μ given by

$$h_\mu \circ \pi_\mu = h \circ i_\mu, \quad \pi_\mu^* \alpha_\mu = i_\mu^* \alpha.$$

- 4 There is a vector field $X_{h_\mu, \alpha_\mu} \in \mathfrak{X}(M_\mu)$ such that

$$\mathbf{T}\pi_\mu \circ X_{h,\alpha} \circ i_\mu = X_{h_\mu, \alpha_\mu} \circ \pi_\mu \quad \text{and} \quad \iota_{X_{h_\mu, \alpha_\mu}} \omega_\mu = dh_\mu + \alpha_\mu.$$

Section 4

Hybrid systems

Hybrid systems

Definition

A **hybrid system** is a 4-tuple $\mathcal{H} = (M, X, S, \Delta)$, formed by

- 1 a manifold M ,
- 2 a vector field $X \in \mathfrak{X}(M)$,
- 3 a submanifold $S \subset M$ of codimension ≥ 1 (**switching surface**),
- 4 an embedding $\Delta: S \rightarrow M$ (**impact map**).

The dynamics generated by \mathcal{H} are the curves $c: I \subseteq \mathbb{R} \rightarrow M$ such that

$$\begin{aligned} \dot{c}(t) &= X(c(t)), & \text{if } c(t) \notin S, \\ c^+(t) &= \Delta(c^-(t)), & \text{if } c(t) \in S, \end{aligned}$$

where

$$c^\pm(t) = \lim_{\tau \rightarrow t^\pm} c(\tau).$$

Hybrid Hamiltonian systems

Definition

A hybrid dynamical system (M, X, S, Δ) is said to be a **hybrid Hamiltonian system** and denoted by \mathcal{H}_h if

- 1 $M \subseteq T^*Q$ is a zero-codimensional submanifold of the cotangent bundle T^*Q of a manifold Q ,
- 2 S projects onto a codimension-one submanifold $\pi_Q(S)$ of Q ,
- 3 $\pi_Q \circ \Delta = \pi_Q$,
- 4 $X = X_h$ is the Hamiltonian vector field of $h \in \mathcal{C}^\infty(T^*Q)$ w.r.t. ω_Q .

A **forced hybrid Hamiltonian system** is defined analogously by replacing X_h with $X_{h,\alpha}$.

Hybrid Hamiltonian systems

Physically,

- Q represents the space of positions,
- T^*Q the phase space,
- X_h the dynamics between the impacts,
- $\pi_Q(S)$ the hypersurface where impacts occur, and
- Δ the change of momenta on the impacts.

Example (The circular billiard)

- Consider a particle in the plane which moves freely inside the surface confined by the unit circle.
- The Hamiltonian is $H = \frac{p_x^2}{2} + \frac{p_y^2}{2}$.
- The switching surface is

$$S = \left\{ (x, y, p_x, p_y) \in T^*\mathbb{R}^2 \mid x^2 + y^2 = 1 \text{ and } (p_x, p_y) \cdot (x, y) > 0 \right\}.$$

- The condition $(p_x, p_y) \cdot (x, y) > 0$ just means that, for an impact to occur, the momenta pointing to the wall must be positive.
- The impact map is $\Delta(x, y, p_x^-, p_y^-) \mapsto (x, y, p_x^+, p_y^+)$, where

$$p_x^+ = p_x^- - 2(xp_x^- + yp_y^-)x,$$

$$p_y^+ = p_y^- - 2(xp_x^- + yp_y^-)y.$$

Hybrid Lie group action

Definition

A Lie group action $\Phi: G \times Q \rightarrow Q$ is called a **hybrid action for \mathcal{H}_h** if its cotangent lift $\Phi^{T^*}: G \times T^*Q \rightarrow T^*Q$ satisfies the following conditions:

- ① h is Φ^{T^*} -invariant, namely, $h \circ \Phi_g^{T^*} = h$ for all $g \in G$,
- ② the restriction $\Phi^{T^*}|_{G \times S}$ is a Lie group action of G on S ,
- ③ the impact map is equivariant w.r.t. this action, i.e.,

$$\Delta \circ \Phi_g^{T^*}|_S = \Phi_g^{T^*} \circ \Delta, \quad \forall g \in G.$$

Hybrid momentum map

Definition

Let $\Phi: G \times Q \rightarrow Q$ be a hybrid action for \mathcal{H}_h . A momentum map $\mathbf{J}: T^*Q \rightarrow \mathfrak{g}^*$ for the cotangent lift action Φ^{T^*} is called a **generalized hybrid momentum map** if, for each connected component $C \subseteq S$ and for each regular value μ_- of \mathbf{J} , there is another regular value μ_+ such that

$$\Delta(\mathbf{J}|_C^{-1}(\mu_-)) \subset \mathbf{J}^{-1}(\mu_+).$$

In particular, if $\mu_- = \mu_+$ it is called a **hybrid momentum map**. A **hybrid regular value** of \mathbf{J} is a regular value of both \mathbf{J} and $\mathbf{J}|_S$.

Hybrid momentum map

In other words, \mathbf{J} is a generalized hybrid momentum map if, for every point in the connected component C of the switching surface S such that the momentum before the impact takes a value of μ_- , the momentum will take a value μ_+ after the impact.

It is a hybrid momentum map if its value does not change with the impacts.

Hybrid reduction

Proposition

If μ_- and μ_+ are regular values of \mathbf{J} such that $\Delta \left(\mathbf{J}|_S^{-1}(\mu_-) \right) \subset \mathbf{J}^{-1}(\mu_+)$, then the isotropy subgroups in μ_- and μ_+ coincide, that is, $G_{\mu_-} = G_{\mu_+}$.

Hybrid reduction

Theorem (Colombo, de León, Eyrea Irazú, L. G., 2022)

Let $\Phi: G \times Q \rightarrow Q$ be a hybrid action on \mathcal{H}_h . Assume that G is connected and that $\Phi^{T^*}: G \times T^*Q \rightarrow T^*Q$ is free and proper. Consider a sequence $\{\mu_i\}_{i \in \mathbb{I} \subseteq \mathbb{N}}$ of hybrid regular values of \mathbf{J} , such that

$\Delta \left(\mathbf{J}|_S^{-1}(\mu_i) \right) \subset \mathbf{J}^{-1}(\mu_{i+1})$. Let $G_{\mu_i} = G_{\mu_0}$ be the isotropy subgroup in μ_i under the co-adjoint action. Then, the reduction leads to a sequence of reduced hybrid forced Hamiltonian systems

$$\mathcal{H}_h^{\mu_i} = \left(\mathbf{J}^{-1}(\mu_i)/G_{\mu_0}, X_{h_{\mu_i}}, \mathbf{J}|_S^{-1}(\mu_i)/G_{\mu_0}, (\Delta)_{\mu_i} \right).$$

Hybrid reduction

$$\begin{array}{ccccccc}
 \dots & \longrightarrow & \mathbf{J}^{-1}(\mu_i) & \longleftarrow & \mathbf{J}|_S^{-1}(\mu_i) & \xrightarrow{\Delta|_{\mathbf{J}^{-1}(\mu_i)}} & \mathbf{J}^{-1}(\mu_{i+1}) & \longleftarrow & \dots \\
 & & \downarrow & & \downarrow & & \downarrow & & \\
 \dots & \longrightarrow & \frac{\mathbf{J}^{-1}(\mu_i)}{G_{\mu_0}} & \longleftarrow & \mathbf{J}|_S^{-1}(\mu_i)/G_{\mu_0} & \xrightarrow{(\Delta)_{\mu_i}} & \frac{\mathbf{J}^{-1}(\mu_{i+1})}{G_{\mu_0}} & \longleftarrow & \dots
 \end{array}$$

Integrable hybrid Hamiltonian systems

- A particular case is when we have the Abelian Lie group action $\Phi: \mathbb{R}^n \times T^*Q \rightarrow T^*Q$ generated by the Hamiltonian flows of n functions f_1, \dots, f_n in involution.
- In that case, we can identify the momentum map with $F = (f_1, \dots, f_n): T^*Q \rightarrow \mathbb{R}^n$.
- We may obtain action-angle coordinates for each time interval between impacts. The action-angle coordinates before and after the impact will be related by Δ .

Section 5

Other results from this thesis

Hamilton–Jacobi theory

- Let $\pi: E \rightarrow B$ is a vector bundle.
- Consider a dynamical system characterized by $X \in \mathfrak{X}(E)$.
- Idea: obtain a section $\gamma \in \Gamma(E)$ such that the following diagram commutes:

$$\begin{array}{ccc}
 E & \xrightarrow{X} & TE \\
 \left. \begin{array}{c} \uparrow \\ \downarrow \end{array} \right\} \gamma & \begin{array}{c} \downarrow \pi \\ \uparrow \end{array} & \left. \begin{array}{c} \uparrow \\ \downarrow \end{array} \right\} T\gamma \\
 B & \xrightarrow{X^\gamma} & TB
 \end{array}$$

- If $c: I \subseteq \mathbb{R} \rightarrow B$ is an integral curve of X^γ , then $\gamma \circ c$ is an integral curve of X .

Hamilton–Jacobi theory


Under certain assumptions on X and γ , the diagram above is commutative iff a PDE known as the **Hamilton–Jacobi (HJ) equation** holds:

- If the bundle is $\pi_Q: T^*Q \rightarrow Q$, the vector field is a forced Hamiltonian vector field $X = X_{h,\alpha}$, and γ is a closed one-form, the HJ equation is

$$\gamma^*(dH + \alpha) = 0.$$

Hamilton–Jacobi theory

- In this dissertation we have also obtained two different HJ equations for a cocontact Hamiltonian vector field $X = X_h$ on $E = \mathbb{R} \times T^*Q \times \mathbb{R}$.
- In that case, one can consider two possible bundles:



Hamilton–Jacobi theory

- We have also studied the HJ theory for hybrid systems.
- Essentially, in that case one has the usual HJ equation for the continuous dynamics between impacts.
- One has to impose a compatibility condition of the form

$$\text{Im}(\Delta \circ \gamma_i) \subset \text{Im} \gamma_{i+1},$$

where γ_i is the solution of the HJ equation between the i -th and $(i + 1)$ -th impacts.

Hamilton–Jacobi theory

- Discrete HJ equations can be obtained by replacing X and X^γ with their discrete flows.
- We have obtained a discrete HJ equation for forced discrete Hamiltonian systems.

Contact Lagrangian systems with impulsive constraints

- Constraints (both holonomic and nonholonomic) with discontinuities can lead to instantaneous changes on dynamical systems.
- Hence, this type of constraints, called **impulsive constraints**, can also be employed to model systems with impacts.
- For instance, one can think of a wall as a constraint.
- Impulsive constraints have been deeply studied in classical mechanics and were given a geometric interpretation in the 1990s by Lacomba and Tulczyjew, Ibort *et al.*, and Cortés and Vinogradov.

Contact Lagrangian systems with impulsive constraints

- In this dissertation, we have extended the theory of impulsive constraints to contact Lagrangian systems.
- In addition, we have proven a **Carnot theorem** for contact Lagrangian systems subject to impulsive constraints, characterizing the changes of energy due to both the contact-type dissipation and the impulsive forces.

Nonsmooth Herglotz variational principle

- Let L be an **action-dependent** Lagrangian function.
- Recall that, roughly speaking, the Herglotz functional \mathcal{A} is given by the ODE

$$\frac{d}{dt}\mathcal{A}[q(t)] = L\left(q(t), \dot{q}(t), \mathcal{A}[q(t)]\right).$$

- One seeks for curves $q: I \subseteq \mathbb{R} \rightarrow Q$ that are extremals of \mathcal{A} .
- Usually, these curves are assumed to be at least \mathcal{C}^2 .
- By considering curves that are \mathcal{C}^0 and piecewise \mathcal{C}^2 we can obtain a variational principle for systems with impacts. The impacts are precisely the points where the curve is not smooth.

Section 6

Future work

Future research

- The Liouville–Arnol'd theorem is a first step in the study of completely integrable contact systems.
- Magri *et al.* studied the relation between bi-Hamiltonian structures, Poisson–Nijenhuis structures and integrable systems. It seems that Jacobi–Nijenhuis structures should have an analogous relation with integrable contact systems.
- We would like to find an algorithm for computing action-angle coordinates in an efficient manner. Perhaps, they are related with solutions of the HJ equation.
- It is pending to consider completely integrable contact systems with critical points, i.e., non-regular values of (f_α) .

Future research

- Other structures employed in the study of classical integrable systems could be generalized to completely integrable contact systems: momentum polytopes, Haantjes tensors, etc.
- We intend to develop a Kolmogorov–Arnol'd–Moser (KAM) theory for contact Hamiltonian systems.
- We would like to study hybrid systems experiencing Zeno effect, i.e., the set of impacts is not discrete.
- It is pending to explore the applicability of our results concerning hybrid systems for mathematical billiard theory.

Publications derived from this thesis

- [1] L. Colombo, M. de León, M. E. Eyrea Irazú, and A. López-Gordón. *Hamilton-Jacobi theory for nonholonomic and forced hybrid mechanical systems*. Accepted on *Geom. Mech.* arXiv: 2211.06252.
- [2] L. Colombo, M. de León, M. Lainz, and A. López-Gordón. *Liouville-Arnold Theorem for Contact Hamiltonian Systems*. 2023. arXiv: 2302.12061.
- [3] L. J. Colombo, M. de León, M. E. Eyrea Irazú, and A. López-Gordón. *Generalized Hybrid Momentum Maps and Reduction by Symmetries of Forced Mechanical Systems with Inelastic Collisions*. 2022. arXiv: 2112.02573.
- [4] L. J. Colombo, M. de León, and A. López-Gordón. “Contact Lagrangian Systems Subject to Impulsive Constraints”. *J. Phys. A: Math. Theor.*, **55**(42) (2022).
- [5] M. de León, M. Lainz, and A. López-Gordón. “Symmetries, Constants of the Motion, and Reduction of Mechanical Systems with External Forces”. *J. Math. Phys.*, **62**(4), p. 042901 (2021).
- [6] M. de León, M. Lainz, and A. López-Gordón. “Discrete Hamilton-Jacobi Theory for Systems with External Forces”. *J. Phys. A: Math. Theor.*, **55**(20) (2022).
- [7] M. de León, M. Lainz, and A. López-Gordón. “Geometric Hamilton–Jacobi Theory for Systems with External Forces”. *J. Math. Phys.*, **63**(2), p. 022901 (2022).

Publications derived from this thesis

- [8] M. de León, M. Lainz, A. López-Gordón, and X. Rivas. “Hamilton–Jacobi Theory and Integrability for Autonomous and Non-Autonomous Contact Systems”. *J. Geom. Phys.*, **187**, p. 104787 (2023).
- [9] M. E. Eyrea Irazú, A. López-Gordón, M. de León, and L. J. Colombo. “Hybrid Routhian Reduction for Simple Hybrid Forced Lagrangian Systems”. In: *2022 European Control Conference (ECC)*. 2022, pp. 345–350.
- [10] J. Gaset, A. López-Gordón, and X. Rivas. “Symmetries, Conservation and Dissipation in Time-Dependent Contact Systems”. *Fortschritte der Phys.*, **71**(8-9), p. 2300048 (2023).
- [11] A. López-Gordón, L. Colombo, and M. de León. “Nonsmooth Herglotz Variational Principle”. In: *2023 American Control Conference (ACC)*. 2023, pp. 3376–3381.
- [12] A. López-Gordón and L. J. Colombo. *On the Integrability of Hybrid Hamiltonian Systems*. Accepted on the Proceedings of the *8th IFAC Workshop on Lagrangian and Hamiltonian Methods for Non Linear Control*. 2023. arXiv: 2312.12152.

Thanks for your kind attention!